

ANNIHILATOR NILPOTENCY OF SKEW BRACES WITH APPLICATIONS TO TORSION AND GENERATING SETS OF SKEW BRACES

(JOINT W. ERIC JESPERS, LEANDRO VENDRAMIN)

Arne Van Antwerpen


THE YANG-BAXTER EQUATION: A PICTURE

Definition

A set-theoretic solution to the Yang-Baxter equation is a tuple (X, r) , where X is a set and $r : X \times X \rightarrow X \times X$ a function such that (on X^3)

$$(r \times \text{id}_X) (\text{id}_X \times r) (r \times \text{id}_X) = (\text{id}_X \times r) (r \times \text{id}_X) (\text{id}_X \times r).$$

For further reference, denote $r(x, y) = (\lambda_x(y), \rho_y(x))$.



DEFINITIONS AND EXAMPLES

Definition

A set-theoretic solution (X, r) is called

- ▶ left (resp. right) non-degenerate, if λ_x (resp. ρ_y) is bijective,
- ▶ non-degenerate, if it is both left and right non-degenerate,
- ▶ involutive, if $r^2 = \text{id}_{X \times X}$,

Examples

- ▶ Twist solution: $r(x, y) = (y, x)$,
- ▶ Lyubashenko, where $f, g : X \rightarrow X$ are maps with $fg = gf$:
 $r(x, y) = (f(y), g(x))$.

THE STRUCTURE MONOID AND GROUP

Definition

Let (X, r) be a set-theoretic solution of the Yang-Baxter equation. Then the group

$$G(X, r) = \langle x \in X \mid xy = \lambda_x(y)\rho_y(x) \rangle,$$

is called the structure group of (X, r) .

WHAT ARE SKEW LEFT BRACES

Definition (Rump, CJO, GV)

Two groups $(A, +)$ and (A, \circ) form a skew left brace $(A, +, \circ)$, if for any $a, b, c \in A$, it holds that

$$a \circ (b + c) = (a \circ b) - a + (a \circ c),$$

where $-a$ denotes the inverse of a in $(A, +)$.

Moreover, if $(A, +)$ is abelian, then $(A, +, \circ)$ is a left brace

EXAMPLES OF SKEW BRACES

Example

1. Every group $(G, +)$ has the skew left brace structure $(G, +, +)$, these are *trivial skew left braces*.
2. The dihedral group $D_{2n} = \langle a, b \mid a^n = b^2 = 1, bab = a^{-1} \rangle$ has a left brace structure, where $a^i b^j + a^k b^l = a^{i+k+jl} b^{j+l}$ with $j, l \in \{0, 1\}$.
3. Radical rings.

CREATING SOLUTIONS ON SKEW BRACES (1)

Definition (Rump, CJO, GV)

Let $(B, +)$ and (B, \circ) be groups on the same set B such that for any $a, b, c \in B$ it holds that

$$a \circ (b + c) = (a \circ b) - a + (a \circ c).$$

Then $(B, +, \circ)$ is called a skew (left) brace

If $(B, +)$ is abelian, one says that $(B, +, \circ)$ is a left brace.

CREATING SOLUTIONS ON SKEW BRACES (1)

Definition (Rump, CJO, GV)

Let $(B, +)$ and (B, \circ) be groups on the same set B such that for any $a, b, c \in B$ it holds that

$$a \circ (b + c) = (a \circ b) - a + (a \circ c).$$

Then $(B, +, \circ)$ is called a skew (left) brace

If $(B, +)$ is abelian, one says that $(B, +, \circ)$ is a left brace.

Denote for $a, b \in B$, the map $\lambda_a(b) = -a + a \circ b$. Then, $\lambda : (B, \circ) \rightarrow \text{Aut}(B, +) : a \mapsto \lambda_a$ is a well-defined group morphism.

CREATING SOLUTIONS ON SKEW BRACES (2)

Theorem

Let $(B, +, \circ)$ be a skew left brace. Denote for any $a, b \in B$, the map $r_B(a, b) = (\lambda_a(b), \overline{\bar{a} + b}) \circ b$. Then (B, r_B) is a bijective non-degenerate solution. Moreover, if $(B, +)$ is abelian, then (B, r_B) is involutive.

Remark

Let (X, r) be a bijective non-degenerate set-theoretic solution. Then, $G(X, r)$ is a skew left brace and carries an associated solution as a skew brace.

THE *-OPERATION IN SKEW LEFT BRACES

Definition

Let $(A, +, \circ)$ be a skew left brace. For any $a, b \in A$, denote

$$a * b = -a + a \circ b - b = \lambda_a(b) - b.$$

Denote $X * Y$ for the additive subgroup generated by $x * y$, where $x \in X, y \in Y$ and $X, Y \subseteq A$.

Example

1. For $(G, +, +)$, one sees that $a * b = 0$. Actually a characterization.
2. For $(D_{2n}, +, \cdot)$ one can see that $(a^i b^j) * (a^k b^l) \in \langle a \rangle$.

SOLUTIONS LIKE LYUBASHENKO'S

Definition (Retraction)

Let (X, r) be a finite bijective non-degenerate set-theoretic solution. Define the relation $x \sim y$ on X , when $\lambda_x = \lambda_y$ and $\rho_x = \rho_y$. Then, there exists a natural set-theoretic solution on X/\sim called the retraction $\text{Ret}(X, r)$.

SOLUTIONS LIKE LYUBASHENKO'S

Definition (Retraction)

Let (X, r) be a finite bijective non-degenerate set-theoretic solution. Define the relation $x \sim y$ on X , when $\lambda_x = \lambda_y$ and $\rho_x = \rho_y$. Then, there exists a natural set-theoretic solution on X/\sim called the retraction $\text{Ret}(X, r)$.

Denote for $n \geq 2$, $\text{Ret}^n(X, r) = \text{Ret}(\text{Ret}^{n-1}(X, r))$. If there exists a positive integer n such that $|\text{Ret}^n(X, r)| = 1$, then (X, r) is called a multipermutation solution

WHY ARE MULTIPERMUTATION SOLUTIONS INTERESTING

Theorem (CJOBVAGI)

Let (X, r) be a finite involutive non-degenerate set-theoretic solution. The following statements are equivalent,

- ▶ the solution (X, r) is a multipermutation solution,
- ▶ the group $G(X, r)$ is left orderable,
- ▶ the group $G(X, r)$ is diffuse,
- ▶ the group $G(X, r)$ is poly- \mathbb{Z} .

Breaks down for non-involutive solutions, as $G(X, r)$ has torsion in that case!

ALL MULTIPERMUTATION SOLUTIONS

Proposition

Let (X, r) be a multipermutation solution, then the skew brace $G(X, r)$ is of nilpotent type.

So we focus attention on so-called skew braces $(B, +, \circ)$ of **nilpotent type**, i.e. $(B, +)$ is a nilpotent group.

NON-ASSOCIATIVE, SO SIDES MATTER

Left Nilpotent

- ▶ $B^{n+1} = B * B^n$ left ideals
- ▶ $|B^k| = 1$, then left nilpotent
- ▶ Nilpotent type:
(B, \circ) nilpotent
- ▶ Example: (C_{2^n}, D_{2^n})

Right nilpotent

- ▶ $B^{(n+1)} = B^{(n)} * B$ ideals
- ▶ $|B^{(k)}| = 1$, then right nilpotent
- ▶ Nilpotent type:
(B, r_B) multipermutation
- ▶ Example: (C_{2n}, D_{2n})

MEASURING MULTIPERMUTATION

Definition

A skew brace $(B, +, \circ)$ is said to be multipermutation, if (B, r_B) is multipermutation.

Equivalently:

- ▶ B is right nilpotent of nilpotent type,
- ▶ The chain $\text{Soc}^n(B)$ ends in B .

Here, $\text{Soc}^{n+1}(B)$ is the pullback in B of $\text{Soc}(B/\text{Soc}^n(B))$ with

$$\text{Soc}(A) = \ker \lambda \cap Z(B, +).$$

CENTRAL NILPOTENCY

Definition

Let B be a skew brace. Denote $Ann(B) = Soc(B) \cap Z(B, \circ)$.

Equivalently,

$$Ann(B) = \{x \in Z(B, +) \mid \lambda_x = id_B, \lambda_y(x) = x \text{ for all } y \in B\}.$$

Definition

Let B be a skew brace. One says that B is centrally nilpotent, if the chain $Ann^n(B)$ ends in B , where $Ann^{k+1}(B)$ is pullback of $Ann(B/Ann^k(B))$.

DESCENDING SERIES

We have an ascending ideal series, what about descending?

$$\Gamma_{n+1}(B, I) = \langle B * \Gamma_n(B, I), \Gamma_n(B, I) * B, [\Gamma_n(B, I), B]_+ \rangle$$

is an ideal in B , if I is an ideal.

Proposition (Bonatto, Jedlicka)

Let B be a skew brace. Then, B is centrally nilpotent, if for some positive integer n we have $\Gamma_n(B, B) = 1$.

STRONGLY NILPOTENT

$$B^{[n]} = \langle B^{[i]} * B^{[n-i]} \mid 1 \leq i \leq n \rangle.$$

Proposition (Smoktunowicz)

Let B be a skew brace. Then, B is strongly nilpotent if and only if B is left and right nilpotent and (B, \circ) is nilpotent.

What if we account for additive commutator?

$$\Gamma_{[n]}(B) = \langle \Gamma_{[i]}(B) * \Gamma_{[n-i]}(B), [\Gamma_{[i]}(B), \Gamma_{[n-i]}(B)]_+ \rangle$$

Proposition (Jespers, AVA, Vendramin)

Let B be a skew brace of nilpotent type. If B is centrally nilpotent, then B is strongly centrally nilpotent. Moreover, B is strongly nilpotent.

NILPOTENCY CLASS

Both the chains $\Gamma_n(B)$ and $\Gamma_{[n]}(B)$ allow to define a notion of nilpotency class of B .

Problem

- ▶ *Can we relate the above nilpotency classes?*
- ▶ *Are there bounds using the additive/multiplicative nilpotency class?*

FINITELY GENERATED

Proposition (Jespers, AVA, Vendramin)

Let B be a centrally nilpotent skew brace with ACC on sub skew braces. TFAE

- ▶ *B is finitely generated as a brace,*
- ▶ *$(B, +)$ is finitely generated as a group,*
- ▶ *(B, \circ) is finitely generated as a group.*

Vice versa, every finitely generated Centrally nilpotent skew brace has ACC on sub skew braces.

TORSION

What is torsion in a skew brace?

Proposition (Jespers, AVA, Vendramin)

Let B be a centrally nilpotent skew brace. Then $T_+(B) = T_0(B)$, which is an ideal of B . Finite, if B is finitely generated.

Proposition (Jespers, AVA, Vendramin)

Let B be a centrally nilpotent skew brace. If $T_+(B) = 0$. Then, $a^n = b^n$ or $na = nb$ implies $a = b$.

REFERENCES

1. E. Jespers, A. Van Antwerpen, L. Vendramin, *Nilpotency of skew braces and multipermutation solutions of the Yang-Baxter equation*, Comm. Cont. Math., to appear.
2. F. Cedó, A. Smoktunowicz, L. Vendramin, *Skew left braces of nilpotent type*, Proc.Lond.Math.Soc. 118(6), 2019.
3. A. Smoktunowicz, *On Engel groups, nilpotent groups, rings, braces and the Yang-Baxter equation*, Trans.Amer.Math.Soc.
4. M. Bonatto, P. Jedlička, *Central nilpotency of skew braces*, J. Alg. Appl.