

Advances in Group Theory and Applications
© 2025 AGTA - www.advgrouptheory.com/journal
22 (2025), pp. 155–175 ISSN: 2499-1287
DOI: 10.32037/agta-2025-018

Characterizing Inner Automorphisms and Realizing Outer Automorphisms

Benjamin Sambale

(*Received May 12, 2024; Accepted Sept. 26, 2024 — Communicated by P. Shumyatsky*)

Abstract

We give elementary proofs of the following two theorems on automorphisms of a finite group G:

(1) An automorphism of G is inner if and only if it extends to an automorphism of every finite group containing G.

(2) There exists a finite group, whose outer automorphism group is isomorphic to G.

The first theorem was proved by Pettet using a graph-theoretical construction given by Heineken–Liebeck. A Lie-theoretical proof of the second theorem was sketched by Cornulier in a *MathOverflow* post. Our proofs are purely group-theoretical.

Mathematics Subject Classification (2020): 20D45, 20F05, 20F28

Keywords: inner automorphism; outer automorphism

1 Introduction

An automorphism α of a group G is called *inner* if there exists some $g \in G$ such that $\alpha(x) = gxg^{-1}$, for all $x \in G$. If G is a subgroup of a group H, it is clear that g still induces an (inner) automorphism of H. In 1987, Schupp [27] has shown conversely that inner automorphisms are characterized by this property,

i.e., if $\alpha \in \mathrm{Aut}(G)$ extends to every group containing G, then α is inner. According to [8], this has answered a question of Macintyre. The question was asked again much later by Bergman, see [1, p. 93], who obtained a partial answer in the language of category theory. Using free products and small cancellation theory, Schupp constructs for a non-inner $\alpha \in \mathrm{Aut}(G)$ an infinite group H such that α does not extend to H (if G is countable, the construction is already contained in the paper by Miller and Schupp [23]). One may asks whether inner automorphisms of *finite* groups G are characterized by the property that they extend to all *finite* groups containing G.

The first step in this direction was made in 1974 by Heineken and Liebeck [13], who constructed a finite p-group P such that the image of the canonical map $\mathrm{Aut}(P) \to \mathrm{Aut}(P/Z(P))$ is isomorphic to G. Their construction relies on a variation of Frucht's theorem on the automorphism group of graphs, and requires a treatment of special cases. Using more advanced graph theoretical theorems, Lawton [20] came up with a shorter proof. Later, Webb [28] has refined the construction (again at the cost of more graph theory) to obtain a special p-group P (that means $P' = Z(P) = \Phi(P)$ is elementary abelian) with the desired property (see also Hughes [14]). Only after Schupp's paper, in 1989, Pettet [24] noted that this result implies Schupp's theorem for finite groups (and more restricted families of groups). In the same paper, he also obtained the dual statement for factor groups (see the theorem below). Eventually, Pettet [25] gave a new proof of Schupp's original theorem with the same graph theoretical approach. An alternative construction of P was established by Bryant and Kovács [4] in 1978 making use of Lie theory (see also the paper by Huppert and Blackburn [16, Theorem VIII.13.5] and the paper by Hartley and Robinson [12]).

The first objective of this paper is to provide a new elementary proof of the following theorem, which avoids small cancellation theory, graph theory and Lie theory.

Theorem 1.1 (Pettet) *For every automorphism α of a finite group G*

the following statements are equivalent:

(1) α *is an inner automorphism.*

(2) α *extends to every finite group containing* G.

(3) α *lifts to every finite group* $\widehat{G}$ *such that* $\widehat{G}/N \simeq G$ *for some characteristic subgroup* $N \leqslant \widehat{G}$.

The strategy is to replace the p-group P in [13] by a semidirect product $N = Q \rtimes P$, where Q is an elementary abelian q-group and P is a p-group of nilpotency class 2. In contrast to the papers cited above, we do not make an effort to minimize N.

Our second goal concerns the inverse problem on automorphism groups. By a theorem of Ledermann and Neumann [21], there exist only finitely many finite groups with a given automorphism group (for an elementary proof see [26]). However, not every group actually occurs as an automorphism group. For instance, it is a popular (and easy) exercise that a nontrivial cyclic group of odd order cannot be an automorphism group. The situation changes if we instead consider the *outer* automorphism group $\mathrm{Out}(H) := \mathrm{Aut}(H)/\mathrm{Inn}(H)$ of groups H. Indeed, Matumoto [22] proved that for every group G there exists a group H such that $\mathrm{Out}(H) \simeq G$ (for finite G this was established earlier by Kojima [19]). Similar to Schupp's paper, the construction of H is based on an HNN-extension and yields infinite groups. In later work, it was shown that H can be chosen to be locally finite, finitely generated (if G is countable), metabelian or simple (see [3], [5], [7] and [11]). Problem 16.59 in the Kourovka Notebook [18] has asked if H can be chosen finite when G is finite. This was answered in 2020 by Cornulier on *MathOverflow* [6].

Theorem 1.2 (Cornulier) *Every finite group is the outer automorphism group of some finite group.*

Its proof uses Lie algebras and is not easy to follow as it is written backwards. In Section 3, we present a purely group-

theoretical proof based on Cornulier's ideas. As in Pettet's theorem, the idea is to construct a semidirect product $P \rtimes Q$, but this time Q is abelian and P is a p-group of exponent p with a certain nilpotency class. Our group is slightly smaller compared to Cornulier's construction.

The construction of our group has already been used by Entin and Tsang [9] to prove that every finite group is isomorphic to the normalizer quotient of a subgroup of a symmetric group.

2 Characterizing inner automorphisms

Our first lemma is a prototype of Theorem 2.4.

Lemma 2.1 *Let G be a group acting faithfully on a cyclic group N. Then every automorphism of $\widehat{G} := N \rtimes G$ normalizing N, centralizes $\widehat{G}/N$.*

Proof — We identify $N = \langle x \rangle$ and G with the natural subgroups of $\widehat{G}$. Let $\alpha \in \mathrm{Aut}(\widehat{G})$ normalizing N. For a fixed $g \in G$ there exist $s, t \in \mathbb{Z}$ with $gxg^{-1} = x^s$ and $\alpha(x) = x^t$. Hence,

$$gx^t g^{-1} = x^{st} = \alpha(x)^s = \alpha(x^s) = \alpha(gxg^{-1}) = \alpha(g)x^t\alpha(g)^{-1}.$$

Since G acts faithfully on $N = \langle x^t \rangle$, we obtain $\alpha(g) \equiv g \pmod{N}$ as desired. □

In the following we develop some elementary facts of finitely presented groups. For elements $x_1, x_2, \ldots$ of a group G we define commutators by $[x_1, x_2] = x_1x_2x_1^{-1}x_2^{-1}$ and

$$[x_1, \ldots, x_k] := [x_1, [x_2, \ldots, x_k]],$$

for $k \geqslant 3$. The commutator subgroup of G is denoted by G'.

Lemma 2.2 *Let p be a prime and $a, b \in \mathbb{Z}$. Then*

$$P := \left\langle x, y \mid [x,x,y] = [y,x,y] = 1,\ x^p = [x,y]^a,\ y^p = [x,y]^b \right\rangle$$

is a non-abelian group of order p^3.

Proof — The commutator relations show that P has nilpotency class $\leqslant 2$, i.e. $P' \leqslant Z(P)$. Hence,

$$[x, y]^p = [x^p, y] = 1$$

and $|P'| \leqslant p$ (see [15, Hilfssatz III.1.3]). It follows that $|P| \leqslant p^3$. To complete the proof, we construct a group of order p^3 realizing the given relations. Suppose first that $p = 2$. In case ab is even, then $P \simeq D_8$ and otherwise $P \simeq Q_8$. Thus, let $p > 2$. If $a \equiv b \equiv 0 \pmod p$, then the extraspecial group of exponent p fulfills the relations as is well-known. Now, let $p \nmid a$ and $a' \in \mathbb{Z}$ such that $aa' \equiv -b \pmod p$. For $y' := x^{a'}y$ we compute

$$(y')^p = x^{pa'}y^p\left[y, x^{a'}\right]^{\binom{p}{2}} = x^{pa'}y^p = [x, y]^{aa'+b} = 1$$

and $[x, y'] = [x, y]$ by [15, Hilfssatz III.1.3]. Hence, replacing y by y', leads to $b = 0$. Besides, this remains true when we replace y by y^{-a}. Then $x^p = [x, y]^{-1} = [y, x]$ and

$$P \simeq \left\langle x, y \mid x^{p^2} = y^p = 1,\ yxy^{-1} = x^{1+p}\right\rangle \simeq C_{p^2} \rtimes C_p.$$

Therefore, the claim follows. □

Lemma 2.3 *Let* F *be the free group in the free generators* $x_1, \ldots, x_n$. *Let* $c_1, \ldots, c_n \in F'$. *For a prime* p, *let* P *be the group generated by* $x_1, \ldots, x_n$ *subject to the relations*

$$x_i^p = c_i \qquad \textit{and} \qquad \left[x_i, x_j, x_k\right] = 1,$$

for all $1 \leqslant i, j, k \leqslant n$. *Then* P/P' *is an elementary abelian* p*-group with basis*

$$\left\{x_iP' : i = 1, \ldots, n\right\}$$

and P' *is an elementary abelian* p*-group with basis*

$$\left\{[x_i, x_j] : 1 \leqslant i < j \leqslant n\right\}.$$

In particular, $|P| = p^{\binom{n+1}{2}}$.

Proof — It follows from $x_i^p = c_i \in P'$ that P/P' is an elementary abelian p-group generated by

$$\{x_iP' : i = 1, \dots, n\}.$$

Since $[x_i, x_j, x_k] = 1$, for all i, j, k, P has nilpotency class $\leqslant 2$. Hence,

$$[xy, z] = [x, z][y, z] \quad \text{and} \quad [x, yz] = [x, y][x, z],$$

for all $x, y, z \in P$ (see [15, Hilfssatz III.1.2]). In particular,

$$[x_i, x_j]^p = [x_i^p, x_j] = [c_i, x_j] = 1.$$

This shows that P' is an elementary abelian p-group generated by $\{[x_i, x_j] : 1 \leqslant i < j \leqslant n\}$.

Now, suppose that $x := \prod_{i<j}[x_i, x_j]^{a_{ij}} = 1$, for some integers $0 \leqslant a_{ij} \leqslant p-1$. We fix $i < j$ and consider the free group F_2 generated by y_i and y_j. Let $\varphi : F \to F_2$ be the homomorphism defined by

$$\varphi(x_k) = \begin{cases} y_k & \text{if } k \in \{i, j\} \\ 1 & \text{otherwise} \end{cases},$$

for $k = 1, \dots, n$. Set

$$P_2 := \langle y_i, y_j \mid [y_i, y_i, y_j] = [y_j, y_i, y_j] = 1,\ y_i^p = \varphi(c_i),\ y_j^p = \varphi(c_j)\rangle.$$

As elements of P_2, $\varphi(c_i)$ and $\varphi(c_j)$ are (possibly trivial) powers of $[y_i, y_j]$. Thus, by Lemma 2.2, P_2 is a non-abelian group of order p^3. Since every relation of P in the elements x_k is satisfied by a relation of P_2 in the elements y_k, φ factors through a homomorphism $\overline{\varphi} : P \to P_2$. It follows that

$$[y_i, y_j]^{a_{ij}} = \overline{\varphi}(x) = 1 \quad \text{and} \quad a_{ij} = 0.$$

This shows that commutators $[x_i, x_j]$ with $i < j$ are linearly independent, so they form a basis of P'. Now, let $x := \prod_{i=1}^{n} x_i^{a_i} \in P'$, for some $0 \leqslant a_i \leqslant p-1$. Then

$$1 = [x_1, x] = \prod_{i=2}^{n} [x_1, x_i]^{a_i} \quad \text{and} \quad a_i = 0,$$

for $i = 2, \ldots, n$. Similarly, we obtain $a_1 = 0$. Therefore, P/P' has rank n as claimed. Finally,

$$|P| = |P'||P/P'| = p^{\binom{n}{2}+n} = p^{\binom{n+1}{2}},$$

which completes the proof. □

Theorem 2.4 *For any finite group* G*, there exist primes* $q > p > |G|$ *and a finite* $\{p, q\}$*-group* N *such that every automorphism of the group* $\widehat{G} := N \rtimes G$ *induces an inner automorphism of* $\widehat{G}/N \simeq G$*.*

Proof — Without loss of generality, we assume that $G \neq 1$. Let $p > |G|$ be a prime, $x_1, \ldots, x_n \in G$ a generating set of G not containing 1 and P the p-group with generators $\{v_g : g \in G\}$ and relations

$$[v_g, v_h, v_k] = 1, \qquad v_g^p = \prod_{i=1}^{n} [v_g, v_{gx_i}]^i,$$

for all $g, h, k \in G$. By Lemma 2.3, P/P' is elementary abelian of rank $|G|$ and P' is elementary abelian with basis $[v_g, v_h]$, where $g < h$, for some fixed total order on G. For a fixed $g \in G$, the elements $\{v_{gh} : h \in G\}$ fulfill the same relations as the v_h. Thus, there exists $\varphi_g \in \mathrm{Aut}(P)$ with

$$\varphi_g(v_h) = v_{gh},$$

for all $h \in G$. This gives rise to a regular action

$$\varphi : G \to \mathrm{Aut}(P), \ g \mapsto \varphi_g.$$

Now, by an elementary special case of Dirichlet's theorem (see,

for example, [10, Theorem 3.1.12]), there exists a prime q such that $p \mid q-1$. Let Q be the elementary abelian q-group with basis $\{w_g : g \in G\}$. Let $w_g \mapsto w_g^\zeta$ be an automorphism of order p of $\langle w_g \rangle$. For $h, g \in G$, define $\gamma_g \in \mathrm{Aut}(Q)$ by

$$\gamma_g(w_h) := \begin{cases} w_g^\zeta & \text{if } h = g \\ w_h & \text{if } h \neq g\,. \end{cases}$$

Let $\gamma : P \to \mathrm{Aut}(Q), v_g \mapsto \gamma_g$ be the homomorphism with kernel P'. This gives rise to the semidirect product $N := Q \rtimes P$ with $Z(N) = P'$. As usual, we identify P and Q with the natural subgroups of N. Then $v_g w_h v_g^{-1} = \gamma_g(w_h)$, for all $g, h \in G$. Again, we have a regular action $\psi : G \to \mathrm{Aut}(Q), g \mapsto \psi_g$ with $\psi_g(w_h) = w_{gh}$. Moreover, φ and ψ are compatible in the sense that

$$\begin{aligned} \varphi_g(v_h)\psi_g(w_k)\varphi_g(v_h)^{-1} &= v_{gh} w_{gk} v_{gh}^{-1} = \gamma_{gh}(w_{gk}) \\ &= \psi_g(\gamma_h(w_k)) = \psi_g(v_h w_k v_h^{-1}), \end{aligned}$$

for all $g, h, k \in G$. In this way, G acts faithfully on N. As before, we identify G and N with subgroups of $\widehat{G} := N \rtimes G$. It follows that

$$g v_h g^{-1} = v_{gh} \quad \text{and} \quad g w_h g^{-1} = w_{gh},$$

for $g, h \in G$.

Now let $\alpha \in \mathrm{Aut}(\widehat{G})$. Since $q > p > |G|$, α normalizes the normal Sylow q-subgroup Q, the normal Hall subgroup N, and in turn $Z(N) = P'$. By the Schur–Zassenhaus theorem, there exists some $y \in N$ such that $\alpha(G) = yGy^{-1}$ (we do not require the Feit–Thompson theorem, because N is solvable). Since y centralizes $\widehat{G}/N$, we may compose α with the inner automorphism induced by y^{-1}. Then α normalizes G. Next, we consider the action of α on

$$N/P' \simeq \bigtimes_{g \in G} \langle w_g, v_g \rangle \simeq (C_q \rtimes C_p)^{|G|}.$$

It follows from

$$\left|C_{N/P'}(\alpha(v_1)P')\right| = \left|C_{N/P'}(v_1P')\right| = p^{|G|}q^{|G|-1}$$

that $\alpha(v_1) \in \langle v_g, w_g\rangle P'$, for some $g \in G$. Composing α with the inner automorphism induced by g^{-1}, we may assume that $g = 1$. Then α induces an automorphism of

$$\langle v_1, w_1\rangle P'/P' \simeq C_q \rtimes C_p.$$

By Lemma 2.1, there exists $t_1 \in P'$ such that

$$\alpha(v_1) \equiv v_1t_1 \pmod{Q}.$$

Moreover, for every $g \in G$ there exists $t_g \in P'$ with

$$\alpha(v_g) = \alpha(gv_1g^{-1}) \equiv \alpha(g)v_1t_1\alpha(g)^{-1} \equiv v_{\alpha(g)}t_g \pmod{Q}.$$

Consequently,

$$\prod_{i=1}^{n}[v_1, v_{x_i}]^i = v_1^p = (v_1t_1)^p \equiv \alpha(v_1)^p \equiv \prod_{i=1}^{n}[v_1t_1, v_{\alpha(x_i)}t_{x_i}]^i$$

$$\equiv \prod_{i=1}^{n}[v_1, v_{\alpha(x_i)}]^i \pmod{Q}.$$

This is a relation in the linearly independent generators $[v_1, v_g]$ of the elementary abelian group P'. Notice that $\alpha(x_i) \neq 1$ and $n < |G| < p$. Comparing exponents reveals $\alpha(x_i) = x_i$, for $i = 1, \ldots, n$. Since $G = \langle x_1, \ldots, x_n\rangle$, α induces the identity automorphism on G. □

How does the group P in the proof of Theorem 2.4 relate to Heineken–Liebeck's construction? Recall that Frucht's theorem states that every finite group G is the automorphism group of some finite graph $\mathcal{G}$. Frucht's graph is based on the Cayley color graph $\mathcal{C}$, which depends on a generating set $x_1, \ldots, x_n$ of G. More precisely, the vertex set of $\mathcal{C}$ is $\{v_g : g \in G\}$ and there is an

arrow $v_g \to v_h$ of color i if and only if $h = gx_i$. Our proof of Theorem 2.4 rests on the elementary fact that the color-preserving automorphism group of $\mathcal{C}$ is isomorphic to G. The purpose of the group Q is to enforce automorphisms to permute the generators of P.

Proof of Theorem 1.1 — The implications (1)⇒(2) and (1)⇒(3) are obvious. Conversely, if α fulfills (2) or (3), then α extends/lifts to the group $\widehat{G}$ constructed in Theorem 2.4. The theorem implies that α is inner. □

Note that, if G is solvable, the proof above shows that (2) and (3) of Pettet's theorem can be restricted to solvable extension groups $\widehat{G}$.

3 Realizing outer automorphisms

We start by proving [6, Lemma 2]. Let $n \in \mathbb{N}$ and $p > n$ be a prime. Let $C_p \rtimes C_{p-1}$ be the holomorph of C_p, i.e. C_{p-1} acts faithfully on C_p. Let S_n be the symmetric group of degree n.

Lemma 3.1 *We have* $\mathrm{Out}((C_p \rtimes C_{p-1})^n) \simeq S_n$.

Proof — The argument is similar as for the group $(C_q \rtimes C_p)^n$ in the proof of Theorem 2.4, although the statement requires the factor $p-1$. Let

$$P = \langle x_1, \dots, x_n \rangle \simeq C_p^n \qquad \text{and} \qquad Q := \langle y_1, \dots, y_n \rangle \simeq C_{p-1}^n$$

such that each y_i induces an automorphism of $\langle x_i \rangle$ of order $p-1$ and centralizes x_j, for $j \neq i$. It is obvious that every permutation $\pi \in S_n$ induces an automorphism α_π of $G := P \rtimes Q$ by permuting the factors $\langle x_i, y_i \rangle$. If $\pi \neq \mathrm{id}$, then α_π is not inner, because it acts non-trivially on the abelian quotient $G/P \simeq Q$. Hence, S_n induces a subgroup of $\mathrm{Out}(G)$.

Conversely, let $\alpha \in \mathrm{Aut}(G)$ be arbitrary. Then α normalizes the normal Sylow p-subgroup P of G. By the Schur–Zassenhaus theorem, we can further assume that $\alpha(Q) = Q$. Since

$$|C_G(\alpha(x_i))| = |C_G(x_i)| = p^n(p-1)^{n-1},$$

there exists some $\pi \in S_n$ with $\alpha(x_i) \in \langle x_{\pi(i)} \rangle$, for $i = 1, \ldots, n$. By composing α with α_π^{-1}, we may assume that $\pi = \mathrm{id}$. A similar argument yields $\alpha(y_i) \in \langle y_i \rangle$. By Lemma 2.1 applied to the group $\langle x_i \rangle \rtimes \langle y_i \rangle$, we have $\alpha(y_i) = y_i$. Since $\mathrm{Aut}(\langle x_i \rangle) \simeq \langle y_i \rangle$, the action of α on $\langle x_i \rangle$ is induced by conjugation with some power of y_i. Composing α with the corresponding inner automorphism, gives $\alpha(x_i) = x_i$ (this will not affect the action of α on $\langle x_j, y_j \rangle$, for $j \neq i$). Doing this for every $i = 1, \ldots, n$, leads to $\alpha = \mathrm{id}$. □

Let F be the free group in the free generators $x_1, \ldots, x_n$, and put

$$F^p = \langle x^p : x \in F \rangle^F \trianglelefteq F$$

be the normal closure of the set of all p-powers in F. Then, the quotient $\overline{F} := F/F^p$ is the free group of rank n and exponent p. We will identify x_i with its image in $\overline{F}$. Define the lower central series by $\overline{F}^{[1]} := \overline{F}$ and

$$\overline{F}^{[k+1]} := \left[\overline{F}, \overline{F}^{[k]}\right] = \left\langle [x, y] : x \in \overline{F},\ y \in \overline{F}^{[k]} \right\rangle$$

for $k \geqslant 1$ as usual. We call

$$\overline{F}_c := \overline{F}/\overline{F}^{[c+1]}$$

the free group of rank n, exponent p and nilpotency class c (we will see in Lemma 3.3 that the class of $\overline{F}_c$ cannot be smaller than c). The adjective "free" is justified by the following universal property: If G is a nilpotent group of exponent p and class $\leqslant c$, and if $y_1, \ldots, y_n \in G$, then there exists a (unique)

homomorphism

$$\overline{F}_c \to G,\ x_i \mapsto y_i,$$

for $i = 1, \dots, n$. We will use this principle frequently in order to verify certain commutator relations in $\overline{F}_c$.

It is easy to show that $\overline{F}^{[k]}/\overline{F}^{[k+1]}$ is generated by the cosets of the k-fold commutators

$$[x_{i_1}, \dots, x_{i_k}]$$

where $1 \leqslant i_1, \dots, i_k \leqslant n$ (see [15, Hilfssatz III.1.11]). It follows that $\overline{F}^{[k]}/\overline{F}^{[k+1]}$ is a finite elementary abelian p-group. In particular, $\overline{F}_c$ is a finite group. The following lemma is certainly known, but I was unable to find a proper reference.

Lemma 3.2 *For every finite group G of exponent p and $k \in \mathbb{N}$, the map*

$$\Phi_k : (G/G')^k \longrightarrow G/G^{[k+1]},$$

$$(g_1G', \dots, g_kG') \longmapsto [g_1, \dots, g_k]G^{[k+1]}$$

is well-defined and multilinear over $\mathbb{F}_p$, i.e. for $1 \leqslant i \leqslant k$, $h \in G$ and $h' \in G'$,

$$[g_1, \dots, g_{i-1}, g_ihh', g_{i+1}, \dots, g_k] \equiv [g_1, \dots, g_k][g_1, \dots, h, \dots, g_k].$$

Proof — If we interpret $[x_1]$ as x_1, then Φ_1 becomes the identity map. Now let $k \geqslant 2$. Recall that

$$[G^{[s]}, G^{[t]}] \leqslant G^{[s+t]}$$

by [15, Hauptsatz III.2.11]. A direct computation shows

$$[xy, z] = [x, y, z][y, z][x, z]$$

and

$$[z, xy] = [xy, z]^{-1} = [z, x][z, y][x, y, z]^{-1},$$

for every $x, y, z \in G$. If $i = 1$, we put $z := [g_2, \dots, g_k] \in G^{[k-1]}$ and obtain

$$\begin{aligned}[g_1hh', g_2, \dots, g_k] &= [g_1h, h', z][h', z][g_1h, z] \equiv [g_1h, z] \\ &\equiv [g_1, h, z][h, z][g_1, z] \equiv [g_1, z][h, z] \\ &\equiv [g_1, \dots, g_k][h, g_2, \dots, g_k] \pmod{G^{[k+1]}}.\end{aligned}$$

It remains to consider the component $i \geqslant 2$. Let

$$z := [g_{i+1}, \dots, g_k].$$

By induction on k, we derive

$$[g_2, \dots, g_ihh', , \dots, g_k] \equiv [g_2, \dots, g_i, z][g_2, \dots, h, z] \pmod{G^{[k]}}$$

and

$$[g_1, \dots, g_ihh', \dots, g_k] \equiv [g_1, \dots, g_i, z][g_1, \dots, h, z] \pmod{G^{[k+1]}},$$

which completes the proof. □

The following result resembles [6, Lemma 5][*].

Lemma 3.3 *Let* $n \geqslant 3$, $\pi \in S_{n-1}$ *and* $0 \leqslant a \leqslant p-1$ *such that*

$$[x_1, \dots, x_{n-1}, x_1] \equiv [x_{\pi(1)}, \dots, x_{\pi(n-1)}, x_{\pi(1)}]^a \pmod{\overline{F}^{[n+1]}}. \quad (3.1)$$

Then $\pi = \mathrm{id}$ *and* $a = 1$.

Proof — By the universal property, it suffices to prove the claim for any elements $x_1, \dots, x_{n-1}$ of a group P with exponent p and nilpotency class $\leqslant n$. Let $P \leqslant GL(n+1, p)$ be the group of upper unitriangular matrices. For $x \in P$ we have

$$x^p - 1 = (x-1)^p = (x-1)^{n+1}(x-1)^{p-n-1} = 0$$

[*] Since the proof of [6, Lemma 5] takes place in the non-nilpotent Lie algebra $\mathfrak{gl}_n$, it is not clear to me that the obtained result can actually be used to prove the main theorem.

since $p > n$. Hence, P has exponent p. For $i < j$, let $E_{ij} \in P$ be the unitriangular matrix with 1 on position (i, j) and zero elsewhere off the diagonal. A direct calculation shows that

$$[E_{ij}, E_{kl}] = E_{il}^{\delta_{jk}} E_{kj}^{-\delta_{il}}, \tag{3.2}$$

where $\delta_{jk}\delta_{il} = 0$ since $i < j$ and $k < l$. An induction shows that $P^{[k]}$ is generated by the matrices E_{ij} with $|j-i| \geqslant k$. In particular, $P^{[n]} = \langle E_{1,n+1} \rangle \simeq C_p$ and $P^{[n+1]} = 1$ (see [15, Satz III.16.3]). So P has indeed nilpotency class n . We define

$$x_1 := E_{12}E_{n,n+1} \qquad \text{and} \qquad x_i := E_{i,i+1},$$

for $i = 2, \dots, n-1$. Then the right hand side of (3.1) is

$$\begin{cases} [x_1, \dots, x_{n-2}, E_{n-1,n+1}] = \dots = [x_1, E_{2,n+1}] = E_{1,n+1} & \text{if } n \geqslant 4 \\ [x_1, E_{24}E_{13}^{-1}] = [E_{12}E_{34}, E_{24}][E_{12}E_{34}, E_{13}]^{-1} = E_{14} & \text{if } n = 3. \end{cases}$$

Suppose first that $\pi(1) \neq 1$. Then x_1 appears only once in

$$c := [x_{\pi(1)}, \dots, x_{\pi(n-1)}, x_{\pi(1)}].$$

By Lemma 3.2, $c \equiv c_1c_2 \pmod{\overline{F}^{[n+1]}}$, where c_1 and c_2 are n-fold commutators in the elements $E_{i,i+1}$. Both c_i contain $x_{\pi(1)}$ twice, so they muss miss some $E_{r,r+1}$. But now each c_i lives inside a direct product of the form

$$Q := \left\{ \begin{pmatrix} Q_1 & 0 \\ 0 & Q_2 \end{pmatrix} : Q_1 \leqslant GL(r,p),\ Q_2 \leqslant GL(n+1-r,p) \right\}.$$

Since Q has nilpotency class $< n$, we derive the contradiction $c \equiv c_1c_2 \equiv 1 \pmod{\overline{F}^{[n+1]}}$.

Therefore, $\pi(1) = 1$. Now, $[x_{\pi(n-1)}, x_1] \neq 1$ implies

$$\pi(n-1) \in \{2, n-1\}$$

by (3.2). Assume first that $\pi(n-1) = n-1$. Then

$$[x_{\pi(n-2)}, x_{n-1}, x_1] = [x_{\pi(n-2)}, E_{n-1,n+1}] \neq 1$$

implies $\pi(n-2) = n-2$. Inductively, one obtains

$$\pi = \mathrm{id},\ c = E_{1,n+1} \quad \text{and} \quad a = 1$$

in this case. Now suppose that $\pi(n-1) = 2$ and without loss of generality, $n \geqslant 4$. Here we use a different realization of $\overline{F}_n$ inside P. More precisely, we reassign $x_i := E_{12}E_{23}\dots E_{n-1,n}$, for $i = 1, \dots, n-2$ and $x_{n-1} := E_{n,n+1}$. Then clearly,

$$c = [\dots, [x_1, x_1]] = 1.$$

On the other hand, the right hand side of (3.1) becomes

$$\begin{aligned}[x_1, \dots, x_{n-1}, x_1] &= [x_1, \dots, x_1, E_{n-1,n+1}]^{-1} \\ &= \dots = [x_1, E_{2,n+1}]^{-1} = E_{1,n+1}^{-1},\end{aligned}$$

a contradiction. □

We have duplicated x_1 in the commutator in Lemma 3.3 to avoid relations of the form

$$[*, \dots, *, x, y] \equiv [*, \dots, *, y, x]^{-1} \pmod{\overline{F}^{[n+1]}}.$$

To prove Cornulier's theorem, let $G = \{g_1, \dots, g_n\}$ be a finite group of order n. We construct a finite group H such that $\mathrm{Out}(H) \simeq G$. Since

$$\mathrm{Out}(1) = 1 \quad \text{and} \quad \mathrm{Out}(C_3) \simeq C_2,$$

we may assume that $n \geqslant 3$ (as in Lemma 3.3). We identify the generators x_i of $\overline{F}$ with x_{g_i} and define

$$N := \left\langle [x_{hg_1}, \dots, x_{hg_{n-1}}, x_{hg_1}] : h \in G \right\rangle \overline{F}^{[n+1]} \leqslant \overline{F}^{[n]}.$$

Since $\overline{F}^{[n]}/\overline{F}^{[n+1]} \leqslant Z(\overline{F}_n)$, it follows that $N \trianglelefteq \overline{F}$. Let

$$P := \overline{F}/N \simeq \overline{F}_n/(N/\overline{F}^{[n+1]}).$$

Notice that P has exponent p and nilpotency class $\leqslant n$. Moreover, $P/P' \simeq \overline{F}/\overline{F}' \simeq C_p^n$. Again we will identify the x_i with their images in P.

Let $\mathbb{F}_p^\times = \langle \zeta \rangle$. For $1 \leqslant i \leqslant n$, the map $x_j \mapsto x_j^{\zeta^{\delta_{ij}}}$ can be extended to an automorphism q_i of $\overline{F}$. By Lemma 3.2,

$$\begin{aligned} q_i([x_{j_1}, \dots, x_{j_n}]) &= [q_i(x_{j_1}), \dots, q_i(x_{j_n})] \\ &\equiv [x_{j_1}, \dots, x_{j_n}]^\gamma \pmod{\overline{F}^{[n+1]}}, \end{aligned}$$

for some $\gamma \in \mathbb{Z}$. In particular, $q_i(N) = N$ and q_i extends to an automorphism of P. Moreover, the group

$$Q := \langle q_1, \dots, q_n \rangle \leqslant \mathrm{Aut}(P)$$

is isomorphic to C_{p-1}^n. Finally, we define $H := P \rtimes Q$. As usual, we regard P and Q as subgroups of H. Then $q_i x_j q_i^{-1} = x_j^{\zeta^{\delta_{ij}}}$, for $1 \leqslant i, j \leqslant n$. Note that $H/P' \simeq (C_p \rtimes C_{p-1})^n$.

The following result implies Cornulier's theorem.

Theorem 3.4 *With the notation above,* $\mathrm{Out}(H) \simeq G$.

Proof — For $h \in G$, the map $x_i \mapsto x_{hg_i}$, $(i = 1, \dots, n)$ can be extended to an automorphism α_h of $\overline{F}$. By the definition of N, we have $\alpha_h(N) = N$. Therefore, we consider α_h as an automorphism of the group P. There exists a similar automorphism $\beta_h \in \mathrm{Aut}(Q)$ with $\beta_h(q_i) = q_{hg_i}$, where q_i is identified with q_{g_i}. Since

$$\begin{aligned} \alpha_h(q_i x_j q_i^{-1}) = \alpha_h(x_j)^{\zeta^{\delta ij}} = x_{hg_j}^{\zeta^{\delta ij}} &= q_{hg_j} x_{hg_i} q_{hg_j}^{-1} \\ &= \beta_h(q_j)\alpha_h(x_i)\beta_h(q_j)^{-1}, \end{aligned}$$

the actions are compatible. This gives rise to a regular action $\alpha : G \to \mathrm{Aut}(H)$. Since $g \neq 1$ acts non-trivially on $H/P \simeq Q$, we obtain that $\alpha(G) \cap \mathrm{Inn}(H) = 1$. Thus, it suffices to show that $\mathrm{Aut}(H) = \alpha(G)\,\mathrm{Inn}(H)$.

To this end, let $\gamma \in \mathrm{Aut}(H)$ be arbitrary. Then γ normalizes the normal Sylow p-subgroup P and P'. By Lemma 3.1, we may assume that γ permutes the factors of H/P'. So there exists a permutation $\pi \in S_n$ such that

$$\gamma(q_i) = q_{\pi(i)} \pmod{P'} \quad \text{and} \quad \gamma(x_i) \equiv x_{\pi(i)} \pmod{P'},$$

for $i = 1, \ldots, n$. This implies

$$[x_{\pi(1)}, \ldots, x_{\pi(n-1)}, x_{\pi(1)}] = \gamma([x_1, \ldots, x_{n-1}, x_1]) = \gamma(1) = 1$$

by Lemma 3.2. This yields an equation inside $N/\overline{F}^{[n+1]}$:

$$\begin{aligned}&[x_{\pi(1)}, \ldots, x_{\pi(n-1)}, x_{\pi(1)}]\\ &\equiv \prod_{h \in G} [x_{hg_1}, \ldots, x_{hg_{n-1}}, x_{hg_1}]^{a_h} \pmod{\overline{F}^{[n+1]}}\end{aligned}$$

for some $0 \leqslant a_h \leqslant p-1$. By the universal property, this equation remains true when we set $x_{\pi(n)} = 1$. For the unique $h \in G$ with $x_{hg_n} = x_{\pi(n)}$, we deduce

$$\begin{aligned}&[x_{\pi(1)}, \ldots, x_{\pi(n-1)}, x_{\pi(1)}]\\ &\equiv [x_{hg_1}, \ldots, x_{hg_{n-1}}, x_{hg_1}]^{a_h} \pmod{\overline{F}^{[n+1]}}.\end{aligned}$$

By Lemma 3.3, $x_{\pi(i)} = x_{hg_i}$, for $i = 1, \ldots, n-1$. Therefore, after composing γ with $\alpha(h)^{-1}$, we may assume that $\pi = 1$. Since $|P'|$ is coprime to $|Q|$ and $\gamma(Q) \leqslant P'Q$, by the Schur–Zassenhaus theorem, there exists $y \in P'$ with $\gamma(Q) = yQy^{-1}$. Since conjugation with y does not affect H/P', we may assume that $\gamma(Q) = Q$. In particular, γ centralizes Q.

Each quotient $P^{[k]}/P^{[k+1]}$ has a basis (as an elementary abelian

group) consisting of some k-fold commutators in the x_i. By concatenation we obtain a basis $c_1, \dots, c_s$ of $\times_{k=1}^{n} P^{[k]}/P^{[k+1]}$. For $c_i \in P^{[k]} \setminus P^{[k+1]}$, we have

$$q_j c_i q_j^{-1} \equiv c_i^{\zeta^l} \pmod{P^{[k+1]}}$$

by Lemma 3.2, where l is the multiplicity of x_j as a component of c_i. Clearly, $l \leqslant k-1 \leqslant n-1 < p-1$. Hence,

$$\zeta^l \equiv 1 \pmod{p}$$

can only hold if $l = 0$. This shows that q_j centralizes c_i if and only if x_j does not appear in c_i. Since the c_i form a basis, it follows that

$$C_P(q_j) = \langle x_i : i \neq j \rangle.$$

On the other hand,

$$q_j \gamma(x_i) q_j^{-1} = \gamma(q_j x_i q_j^{-1}) = \gamma(x_i),$$

for $j \neq i$, shows that

$$\gamma(x_i) \in \bigcap_{j \neq i} C_P(q_j) = \langle x_i \rangle.$$

Since we already know that $\gamma(x_i) \equiv x_i \pmod{P'}$, we conclude that $\gamma(x_i) = x_i$, for $i = 1, \dots, n$ and $\gamma = \mathrm{id}$, as desired. □

In order to estimate $|H|$ in terms of n, we consider the free nilpotent Lie algebra L over $\mathbb{Q}$ of rank n and class n. By Witt's formula,

$$\dim L = \sum_{k=1}^{n} \frac{1}{k} \sum_{d|k} \mu(d) n^{k/d},$$

where μ is the Möbius function. This number grows roughly as n^{n-1} (see [2, Lemma 20.7]). Let us notice that $\mathbb{F}_p \otimes L$ is the corresponding free nilpotent Lie algebra over $\mathbb{F}_p$. In addition, since $p > n$, the Lazard correspondence turns $\mathbb{F}_p \otimes L$

into $\bar{F}_n$ (see [17, Example 10.24]). In particular, $|\bar{F}_n| = p^{\dim L}$. Moreover, an application of Lemma 3.3 reveals that

$$\left|N/\bar{F}^{[n+1]}\right| = p^n \sim (p-1)^n = |Q|.$$

Altogether, the order of magnitude of $|H|$ is $p^{n^{n-1}}$ (the estimate n^n in [6] is unjustified).

As a concrete example, consider $G \simeq C_3$. Here we can take $p = 5$. Then

$$|\bar{F}_3| = 5^{14}, \quad |P| = 5^{11} \quad \text{and} \quad |H| = 2^6 5^{11}.$$

Cornulier's construction yields a group of order $2^6 5^{29}$, as he remarked at the end of [6].

One may ask if the group H can also be used to prove Pettet's theorem. This does not seem to be easy, since it is not clear whether every automorphism of $H \rtimes G$ normalizes H. Conversely, the group N in Theorem 2.4 has outer automorphisms, which do not come from G.

REFERENCES

[1] G.M. Bergman: "An inner automorphism is only an inner automorphism, but an inner endomorphism can be something strange", *Publ. Mat.* 56 (2012), 91–126.

[2] S.R. Blackburn – P.M. Neumann – G. Venkataraman: "Enumeration of Finite Groups", *Cambridge University Press*, Cambridge (2007).

[3] G. Braun – R. Göbel: "Outer automorphisms of locally finite p-groups", *J. Algebra* 264 (2003), 55–67.

[4] R.M. Bryant – L.G. Kovács: "Lie representations and groups of prime power order", *J. London Math. Soc.* (2) 17 (1978), 415–421.

[5] I. Bumagin – D.T. Wise: "Every group is an outer automorphism group of a finitely generated group", *J. Pure Appl. Algebra* 200 (2005), 137–147.

[6] Y. Cornulier: "Is every finite group the outer automorphism group of a finite group?", *MathOverflow*; https://mathoverflow.net/q/372563.

[7] M. Droste – M. Giraudet – R. Göbel: "All groups are outer automorphism groups of simple groups", *J. London Math. Soc.* (2) 64 (2001), 565–575.

[8] M. Dugas – R. Göbel: "Outer automorphisms of groups", *Illinois J. Math.* 35 (1991), 27–46.

[9] A. Entin – C. Tsang: "Normalizer quotients of symmetric groups and inner holomorphs"; *arXiv:2408.07133v1*.

[10] B. Fine – G. Rosenberger: "Number Theory", *Birkhäuser/Springer*, Cham (2016).

[11] R. Göbel – A.T. Paras: "Outer automorphism groups of metabelian groups", *J. Pure Appl. Algebra* 149 (2000), 251–266.

[12] B. Hartley – D.J.S. Robinson: "On finite complete groups", *Arch. Math.* (*Basel*) 35 (1980), 67–74.

[13] H. Heineken – H. Liebeck: "The occurrence of finite groups in the automorphism group of nilpotent groups of class 2", *Arch. Math.* (*Basel*) 25 (1974), 8–16.

[14] A. Hughes: "Automorphisms of nilpotent groups and supersolvable orders", *Proc. Sympos. Pure Math.* 37 (1980), 205–207.

[15] B. Huppert: "Endliche Gruppen I", *Springer*, Berlin (1967).

[16] B. Huppert – N. Blackburn: "Finite Groups II", *Springer*, Berlin (1982).

[17] E.I. Khukhro: "p-Automorphisms of Finite p-Groups", *Cambridge University Press*, Cambridge (1998).

[18] E.I. Khukhro – V.D. Mazurov: "The Kourovka Notebook. Unsolved Problems in Group Theory", 18th edition, *Russian Academy of Sciences Siberian Division, Institute of Mathematics*, Novosibirsk (2014).

[19] S. Kojima: "Isometry transformations of hyperbolic 3-manifolds", *Topology Appl.* 29 (1988), 297–307.

[20] R. Lawton: "A note on a theorem of Heineken and Liebeck", *Arch. Math.* (*Basel*) 31 (1978/79), 520–523.

[21] W. Ledermann – B.H. Neumann: "On the order of the automorphism group of a finite group I", *Proc. Roy. Soc. London. Ser. A.* 233 (1956), 494–506.

[22] T. Matumoto: "Any group is represented by an outerautomorphism group", *Hiroshima Math. J.* 19 (1989), 209–219.

[23] C.F. Miller – P.E. Schupp: "Embeddings into Hopfian groups", *J. Algebra* 17 (1971), 171–176.

[24] M.R. Pettet: "On inner automorphisms of finite groups", *Proc. Amer. Math. Soc.* 106 (1989), 87–90.

[25] M.R. Pettet: "Characterizing inner automorphisms of groups", *Arch. Math.* (*Basel*) 55 (1990), 422–428.

[26] B. Sambale: "On a theorem of Ledermann and Neumann", *Amer. Math. Monthly* 127 (2020), 827–834.

[27] P.E. Schupp: "A characterization of inner automorphisms", *Proc. Amer. Math. Soc.* 101 (1987), 226–228.

[28] U.H.M. Webb: "The occurrence of groups as automorphisms of nilpotent p-groups", *Arch. Math.* (*Basel*) 37 (1981), 481–498.

Benjamin Sambale
Institut für Algebra, Zahlentheorie und Diskrete Mathematik
Leibniz Universität Hannover
Welfengarten 1, 30167 Hannover, Germany
e-mail: sambale@math.uni-hannover.de